

Algorithms in Genome Research
Winter 2020/2021

Exercises

Number 7, Discussion: 2021 January 08

1. Given a list of peaks from a Tandem Mass Spectrum (MS/MS) for peptide de-novo sequencing, one important obstacle for recovering the peptide sequence is to assign peaks to the main ion types b and y (prefix and suffix strings of the peptide sequence).

If we know that only b -ions are present in the spectrum, then recovering the sequence becomes simple. Describe an algorithm to do so: Input are the parent mass W and an ordered list of prefix masses w_1, \dots, w_k . (Some peaks may be missing, though.)

2. We modify the above problem such that there are “noise peaks” (of unknown origin) in the mass spectrum. Describe an algorithm that finds a peptide sequence maximizing the number of explained peaks. The algorithm should run in $O(k|\Sigma|)$ time where k is the number of peaks and Σ is the underlying alphabet of amino acids. Hint: use dynamic programming.

3. Word puzzling:

- How many different words can be built by using all the letters of the word GLÜHWEIN exactly once? Compute the actual value.
- How many different words can be built by using all the letters of the word TEELICHT exactly once? Note that all words need to have the same length and must use the letters the specified number of times: For ABA, there are three such words, AAB, ABA, BAA.
- Try and find a general formula. Hint: For GLÜHWEIN, the formula depends only on the length of the word, but not for TEELICHT - what else does it depend on?

4. Suppose that we do not know the order of characters in a string: For example, the strings AACCC, ACACC, \dots , CCCAA are indistinguishable to us. We call such “strings without order” *compomers* (denoted A_2C_3 for our example). The *length* of a compomer is the length of the corresponding string (5 in our example).

- Let $\Sigma = \{A, C, G, T\}$ be our alphabet, then there exist 4 compomers of length 1 (A_1, C_1, G_1, T_1) and 10 compomers of length 2 ($A_2, A_1C_1, A_1G_1, A_1T_1, C_2, C_1G_1, C_1T_1, G_2, G_1T_1, T_2$). How many compomers exist of lengths 3 and 4?
- Derive a general formula for the number of compomers of length n over an arbitrary alphabet Σ of size σ .